



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

## Directed strongly regular graphs with rank 5

Jørgensen, Leif Kjær

*Publication date:*  
2014

*Document Version*  
Early version, also known as pre-print

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*

Jørgensen, L. K. (2014). *Directed strongly regular graphs with rank 5*. Department of Mathematical Sciences, Aalborg University. Research Report Series No. R-2014-05

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### Take down policy

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.

**Directed strongly regular graphs with rank 5**

by

Leif Kjær Jørgensen

R-2014-05

May 2014

DEPARTMENT OF MATHEMATICAL SCIENCES  
AALBORG UNIVERSITY

Fredrik Bajers Vej 7 G ■ DK-9220 Aalborg Øst ■ Denmark

Phone: +45 99 40 99 40 ■ Telefax: +45 99 40 35 48

URL: <http://www.math.aau.dk>



# Directed Strongly Regular Graphs with rank 5

L. K. Jørgensen

Dept. of Mathematical Sciences

Aalborg University

Fr. Bajers Vej 7, 9220 Aalborg, Denmark.

leif@math.aau.dk

May 20, 2014

## Abstract

From the parameters  $(n, k, t, \lambda, \mu)$  of a directed strongly regular graph (dsrg) A. Duval (1988) showed how to compute the eigenvalues and multiplicities of the adjacency matrix, and thus the rank of the adjacency matrix. For every rational number  $q$ , where  $\frac{1}{5} \leq q \leq \frac{7}{10}$ , there is feasible (i.e., satisfying Duval's conditions) parameter set for a dsrg with rank 5 and with  $\frac{k}{n} = q$ .

In this paper we show that there exist a dsrg with such a feasible parameter set only if  $\frac{k}{n}$  is  $\frac{1}{5}$ ,  $\frac{1}{3}$ ,  $\frac{2}{5}$ ,  $\frac{1}{2}$ ,  $\frac{3}{5}$ ,  $\frac{2}{3}$ . Every dsrg with rank 5 therefore has parameters of a known graph. The proof is based on an enumeration of  $5 \times 5$  matrices with entries in  $\{0, 1\}$ .

## 1 Introduction

A directed strongly regular graph with parameters  $(n, k, t, \lambda, \mu)$  is a  $k$ -regular directed graph on  $n$  vertices such that every vertex is on  $t$  2-cycles (which may be thought of as undirected edges), and the number of paths of length 2 from a vertex  $x$  to a vertex  $y$  is  $\lambda$  if there is an edge directed from  $x$  to  $y$  and it is  $\mu$  otherwise. Thus the adjacency matrix  $A$  satisfies

$$A^2 = tI + \lambda A + \mu(J - I - A) \quad \text{and} \quad AJ = JA = kJ.$$

It is usually assumed that  $0 < t < k$ . These graphs were introduced by A. Duval [4], who also showed that the spectrum of  $A$  may be computed from the parameters.

In some cases 0 is an eigenvalue of large multiplicity and then the rank of  $A$  is small. In [8], we proved that there exists a dsrg with parameters  $(n, k, t, \lambda, \mu)$  and with adjacency matrix of rank 3 if and only if the parameters are either  $(6m, 2m, m, 0, m)$  or  $(8m, 4m, 3m, m, 3m)$ , for some integer  $m$ , and there exists one with rank 4 if and only if  $(n, k, t, \lambda, \mu)$  is either  $(6m, 3m, 2m, m, 2m)$  and  $(12m, 3m, m, 0, m)$ , for an integer  $m$ . For rank 3, this was proved independently in [6].

The main theorem in this paper is a characterization of parameters with rank 5.

**Theorem 1** *There exists a directed strongly regular graph with parameters  $(n, k, t, \lambda, \mu)$  and with adjacency matrix of rank 5 if and only if the parameter set is one of the following:  $(20m, 4m, m, 0, m)$ ,  $(36m, 12m, 5m, 2m, 5m)$ ,  $(10m, 4m, 2m, m, 2m)$ ,  $(16m, 8m, 5m, 3m, 5m)$ ,  $(20m, 12m, 9m, 6m, 9m)$ , or  $(18m, 12m, 10m, 7m, 10m)$ , for some positive integer  $m$ .*

Note that in these results we assume that  $t < k$ . A dsrg with  $t = k$ , eigenvalue 0 and with rank  $r$  is an undirected complete  $r$ -partite graph, that exists for every  $r \geq 2$ . In the following table we list the possible values of  $\frac{k}{n}$  for which there exists a dsrg with rank  $r \leq 5$ , including  $\frac{r-1}{r}$  that we get when  $t = k$ . We see that then the list is symmetric around  $\frac{1}{2}$  for each  $r \leq 5$ . It would be interesting to know if this is also true for  $r \geq 6$ .

Rank	Values of $k/n$
2	$\frac{1}{2}$
3	$\frac{1}{3}, \frac{1}{2}, \frac{2}{3}$
4	$\frac{1}{4}, \frac{1}{2}, \frac{3}{4}$
5	$\frac{1}{5}, \frac{1}{3}, \frac{2}{5}, \frac{1}{2}, \frac{3}{5}, \frac{2}{3}, \frac{4}{5}$

For the proof of the *if* part of Theorem 1 we refer to known constructions. Duval [4] proved that if for one of the six families of parameters sets there exists a dsrg for  $m = 1$  then there exists a dsrg for every parameter set in that

family. This construction replaces each vertex by a set of  $m$  independent vertices, and it works when  $t = \mu$ . Graphs with parameters  $(20, 4, 1, 0, 1)$  and  $(10, 4, 2, 1, 2)$  were also constructed by Duval. A dsrg with parameters  $(20, 7, 4, 3, 2)$  was constructed in [10]. Its complement has parameters  $(20, 12, 9, 6, 9)$ . A dsrg with parameters  $(18, 5, 3, 2, 1)$  and the complementary dsrg with parameters  $(18, 12, 10, 7, 10)$  was constructed in [5]. A dsrg with parameters  $(36, 12, 5, 2, 5)$  was constructed in [3]. In [7] we constructed dsrgs with parameters  $(4r - 4, 2r - 3, r - 1, r - 2, r - 2)$ , for every  $r \geq 3$ . The complement has parameters  $(4r - 4, 2r - 2, r, r - 2, r)$  and it has rank  $r$ . This proves that a  $(16, 8, 5, 3, 5)$  exists and that for every  $r \geq 3$  there is a dsrg with rank  $r$  and  $\frac{k}{n} = \frac{1}{2}$ . In several of these cases, there exist (many) non-isomorphic graphs. In fact some constructions of directed strongly regular graphs with low rank involve some degree of randomness, see e.g. [1].

## 2 Preliminaries

Duval [4] proved the following conditions for the existence of a dsrg.

**Theorem 2** *Suppose that there exists a directed strongly regular graph with parameters  $(n, k, \mu, \lambda, t)$ .*

*Then the parameters satisfy*

$$k(k + (\mu - \lambda)) = t + (n - 1)\mu \quad (1)$$

and

$$0 \leq \lambda < t, \quad 0 < \mu \leq t, \quad -2(k - t - 1) \leq \mu - \lambda \leq 2(k - t). \quad (2)$$

*The eigenvalues of the adjacency matrix are*

$$k > \rho = \frac{1}{2}(-(\mu - \lambda) + d) > \sigma = \frac{1}{2}(-(\mu - \lambda) - d),$$

*for some positive integer  $d$ , where  $d^2 = (\mu - \lambda)^2 + 4(t - \mu)$ . The multiplicities are*

$$1, \quad -\frac{k + \sigma(n - 1)}{\rho - \sigma}, \quad \frac{k + \rho(n - 1)}{\rho - \sigma}, \quad (3)$$

*respectively.*

We say that  $(n, k, \mu, \lambda, t)$  is a feasible parameter set if the conditions (1) and (2) are satisfied and the multiplicities in (3) are positive integers, and we say that  $(n, k, \mu, \lambda, t)$  is a realizable parameter set if there exists a directed strongly regular graph with these parameters.

From Theorem 2 it follows that we have an eigenvalue  $\rho = 0$  if and only if  $d = \mu - \lambda$ , i.e.,  $t = \mu$ . If 0 is an eigenvalue of the adjacency matrix then the rank of the adjacency matrix is the sum of multiplicities of non-zero eigenvalues, i.e.,

$$\text{rank} = 1 + \frac{k + \rho(n-1)}{\rho - \sigma} = 1 + \frac{k}{\mu - \lambda} = 1 + \frac{k}{d}.$$

Thus we define the rank of a feasible parameter set with  $t = \mu$  to be  $1 + \frac{k}{\mu - \lambda}$  (even if no directed strongly regular graph exists with these parameters).

If the rank of a dsrg and the value of  $\frac{k}{n}$  is known then we can find the parameters.

**Proposition 3** *If  $(n, k, t, \lambda, \mu)$  are the parameters of a dsrg with rank 5 and with  $\frac{k}{n} = \frac{a}{b}$  where  $a$  and  $b$  are relatively prime integers then*

$$(n, k, t, \lambda, \mu) = \left( \frac{(r-1)b^2}{c}m, \frac{(r-1)ab}{c}m, \frac{ra^2}{c}m, \frac{(ar-b)a}{c}m, \frac{ra^2}{c}m \right),$$

for some positive integer  $m$ , where  $c$  is the greatest common divisor of  $((r-1)b^2, (r-1)ab, ra^2, (ar-b)a, ra^2)$ .

**Proof** Using that  $1 + \frac{k}{d} = r$ ,  $d = \mu - \lambda$ ,  $\mu = t$ ,  $\frac{k}{n} = \frac{a}{b}$  and equation 1, we get  $(n, k, t, \lambda, \mu) = (\frac{b}{a}(r-1)d, (r-1)d, \frac{a}{b}rd, (\frac{a}{b}r-1)d, \frac{a}{b}rd)$ , where  $d$  is an integer satisfying that all parameters are integers. Replacing  $d$  by  $ab\frac{m}{c}$  we get the required result.  $\square$

In [9], we proved from Theorem 2 that for a positive integer  $r$  there exists a feasible parameter set  $(n, k, \mu, \lambda, t)$  with rank  $r$  and with  $\frac{k}{n} = q$  if and only if  $q$  is a rational number in the interval  $[\frac{1}{r}, \frac{2r-3}{2r}]$ .

However, we also proved in [9] that there are only finitely many values of  $\frac{k}{n}$  for which there exists a  $k$ -regular directed graph on  $n$  vertices.

We say that a  $\{0, 1\}$  matrix is  $k$ -regular if it has exactly  $k$  ones in each row and in each column.

**Theorem 4 ([9])** *For any positive integer  $r$  the set of values of  $\frac{k}{n}$ , for which there exists a  $k$ -regular  $n \times n$  matrix of rank  $r$  is finite.*

*In particular, if there is a  $k$ -regular  $n \times n$  matrix of rank 5 then*

$$\frac{k}{n} \in \left\{ \frac{1}{5}, \frac{1}{4}, \frac{2}{7}, \frac{1}{3}, \frac{3}{8}, \frac{2}{5}, \frac{3}{7}, \frac{4}{9}, \frac{1}{2}, \frac{5}{9}, \frac{4}{7}, \frac{3}{5}, \frac{5}{8}, \frac{2}{3}, \frac{5}{7}, \frac{3}{4}, \frac{4}{5} \right\}.$$

The result for rank 5 was based on computer enumeration of all  $5 \times 5$   $\{0, 1\}$ -matrices with rank 5.

The idea of the proof of Theorem 1 is that a large  $k$ -regular matrix with low rank is forced to have many identical rows (Section 3), but the adjacency matrix of a dsrg can not have too many identical rows (Section 4).

### 3 Regular rank 5 matrices

**Lemma 5** *A  $k$ -regular  $n \times n$  matrix with rank 5 and  $\frac{k}{n} = \frac{4}{7}$  or  $\frac{k}{n} = \frac{3}{7}$  has a set of  $\frac{2}{7}n$  identical rows or a set of  $\frac{2}{7}n$  identical columns.*

**Proof** Let  $A$  be a  $k$ -regular  $n \times n$  matrix with rank 5. If  $\frac{k}{n} = \frac{4}{7}$  then  $J - A$  is a  $(n - k)$ -regular matrix with rank 5 and with  $\frac{n-k}{n} = \frac{3}{7}$ . So let us assume that  $\frac{k}{n} = \frac{3}{7}$ .

Let  $M$  be  $5 \times n$  submatrix of  $A$  of rank 5 and let  $C = [\mathbf{c}_1, \mathbf{c}_2, \mathbf{c}_3, \mathbf{c}_4, \mathbf{c}_5]$  be a  $5 \times 5$  submatrix of  $M$  of rank 5 and with columns  $\mathbf{c}_1, \dots, \mathbf{c}_5$ . There exists unique real numbers  $\alpha_1, \dots, \alpha_5$  so that  $\sum_{i=1}^5 \alpha_i \mathbf{c}_i = \mathbf{j}_5$ , the all 1 vector, as  $k\mathbf{j}_5$  is the sum of all columns of  $M$ . If  $\mathbf{a}_1, \dots, \mathbf{a}_5$  are the corresponding columns of  $A$  then  $\sum_{i=1}^5 \alpha_i \mathbf{a}_i = \mathbf{j}_n$ . Taking dot product of this equation with  $\mathbf{j}_n$  shows that  $k \sum_{i=1}^5 \alpha_i = n$ , i.e,  $\alpha_1 + \dots + \alpha_5 = \frac{7}{3}$ .

We say that two matrices are equivalent if one of them can be obtained from the other by permuting rows and permuting columns. Using computer we find that there are four equivalence classes of  $5 \times 5$   $\{0, 1\}$ -matrices of rank 5 satisfying  $\sum_{i=1}^5 \alpha_i \mathbf{c}_i = \mathbf{j}_n$  with  $\alpha_1 + \dots + \alpha_5 = \frac{7}{3}$ . These equivalence classes are represented by

$$C_1 = \begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \end{bmatrix}, C_2 = \begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 & 0 \end{bmatrix}, C_3 = \begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix},$$

and  $C_2^T$ .

Each matrix  $C$  (other than  $C_2^T$ ) representing an equivalence class is chosen so that the binary number  $C_{11} \dots C_{15} C_{21} \dots C_{25} \dots C_{51} \dots C_{55}$  is largest. (We assume that the rows of  $M$  are permuted similarly.)

Any additional column of  $M$  is a linear combination

$$\sum_{i=1}^5 \beta_i \mathbf{c}_i, \quad \text{where } \beta_1 + \dots + \beta_5 = 1. \quad (4)$$

The last equation is to ensure that the sum of entries in the new column of  $A$  is  $k$ . For the matrix  $C = C_3$  this condition is not satisfied by any  $\{0, 1\}$  vector other than  $\mathbf{c}_1, \dots, \mathbf{c}_5$ . Thus every column of  $M$  is identical to one of  $\mathbf{c}_1, \dots, \mathbf{c}_5$ . As  $\frac{1}{3}(\mathbf{c}_1 + \mathbf{c}_2 + \mathbf{c}_3 + \mathbf{c}_4 + 3\mathbf{c}_5) = \mathbf{j}_5$ , column  $\mathbf{c}_5$  appears  $\frac{3}{7}n$  times.

Also if  $C = C_2$  then every column must be identical to a column of  $C$ . As  $\frac{1}{3}(\mathbf{c}_1 + \mathbf{c}_2 + \mathbf{c}_3 + 2\mathbf{c}_4 + 2\mathbf{c}_5) = \mathbf{j}_5$ , columns  $\mathbf{c}_4$  and  $\mathbf{c}_5$  each appear  $\frac{2}{7}n$  times. If  $C = C_2^T$  then  $C_2$  is submatrix  $A^T$ , therefore  $A$  has  $\frac{2}{7}n$  identical rows.

Thus we may assume that every  $5 \times 5$  submatrix of  $A$  of rank 5 is equivalent to  $C_1$ . For  $C = C_1$  the possible  $\{0, 1\}$  vectors satisfying equation 4 are  $\mathbf{c}_1, \dots, \mathbf{c}_5, \mathbf{c}_6 = -\mathbf{c}_2 + \mathbf{c}_3 + \mathbf{c}_4 = (0, 1, 0, 0, 1)^T$  and  $\mathbf{c}_7 = \mathbf{c}_2 - \mathbf{c}_3 + \mathbf{c}_5 = (0, 0, 1, 1, 0)^T$ . We know that the columns  $\mathbf{c}_1, \dots, \mathbf{c}_5$  appear in  $M$ . If  $\mathbf{c}_6$  also appears in  $M$  then  $[\mathbf{c}_1 \ \mathbf{c}_3 \ \mathbf{c}_4 \ \mathbf{c}_5 \ \mathbf{c}_6]$  is equivalent to  $C_2^T$ . If  $\mathbf{c}_7$  appears in  $M$  then  $[\mathbf{c}_1 \ \mathbf{c}_2 \ \mathbf{c}_3 \ \mathbf{c}_4 \ \mathbf{c}_7]$  is equivalent to  $C_2^T$ . Thus every column of  $M$  is identical to one of  $\mathbf{c}_1, \dots, \mathbf{c}_5$ . As  $\frac{1}{3}(\mathbf{c}_1 + \mathbf{c}_2 + \mathbf{c}_3 + 2\mathbf{c}_4 + 2\mathbf{c}_5) = \mathbf{j}_5$ , columns  $\mathbf{c}_4$  and  $\mathbf{c}_5$  each appear  $\frac{2}{7}n$  times in  $M$ .  $\square$

In fact a stronger result follows from the above proof.

**Corollary 6** *Let  $A$  be  $\frac{3}{7}n$ -regular  $n \times n$  matrix of rank 5. Then either  $A$  or  $A^T$  satisfies one of the following properties.*

- *There is a set of  $\frac{3}{7}n$  identical rows, or*
- *There are two disjoint sets of  $\frac{2}{7}n$  identical rows.*

**Lemma 7** *A  $k$ -regular  $n \times n$  matrix with rank 5 and  $\frac{k}{n} = \frac{3}{8}$  or  $\frac{k}{n} = \frac{5}{8}$  has a set of  $\frac{3}{8}n$  identical rows and a set of  $\frac{3}{8}n$  identical columns.*

**Proof** There is only one equivalence class of  $5 \times 5$   $\{0, 1\}$ -matrices  $C = [\mathbf{c}_1 \dots \mathbf{c}_5]$  of rank 5 satisfying  $\sum_{i=1}^5 \alpha_i \mathbf{c}_i = \mathbf{j}_5$  with  $\alpha_1 + \dots + \alpha_5 = \frac{8}{3}$ . This class is represented by the matrix  $C = C_{\frac{3}{8}}$  shown below. The columns  $\mathbf{c}_1 \dots \mathbf{c}_5$



are the only  $\{0, 1\}$ -vectors of the form  $\sum_{i=1}^5 \beta_i \mathbf{c}_i$  with  $\beta_1 + \dots + \beta_5 = 1$ . As  $\frac{1}{3}(\mathbf{c}_1 + \mathbf{c}_2 + \mathbf{c}_3 + 2\mathbf{c}_4 + 3\mathbf{c}_5)$ , any  $\frac{3}{8}n$ -regular  $n \times n$  matrix with rank 5 has a set of  $\frac{3}{8}n$  identical columns. As  $C$  is equivalent to its transpose, the same holds for rows.  $\square$

$$C_{\frac{3}{8}} = \begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}, C_{\frac{2}{7}} = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}, C_{\frac{4}{9}} = \begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 \end{bmatrix}.$$

The cases  $\frac{2}{7}$  and  $\frac{4}{9}$  have similar proofs.

**Lemma 8** *A  $k$ -regular  $n \times n$  matrix with rank 5 and  $\frac{k}{n} = \frac{2}{7}$  or  $\frac{k}{n} = \frac{5}{7}$  has a set of  $\frac{2}{7}n$  identical rows and a set of  $\frac{2}{7}n$  identical columns.*

**Lemma 9** *A  $k$ -regular  $n \times n$  matrix with rank 5 and  $\frac{k}{n} = \frac{4}{9}$  or  $\frac{k}{n} = \frac{5}{9}$  has a set of  $\frac{1}{3}n$  identical rows and a set of  $\frac{1}{3}n$  identical columns.*

**Lemma 10** *A  $k$ -regular  $n \times n$  matrix with rank 5 and  $\frac{k}{n} = \frac{1}{4}$  or  $\frac{k}{n} = \frac{3}{4}$  has a set of  $\frac{1}{4}n$  identical rows or a set of  $\frac{1}{4}n$  identical columns.*

**Proof** Let  $C = [\mathbf{c}_1 \dots \mathbf{c}_5]$  be a full rank  $5 \times 5$  submatrix of a  $\frac{n}{4}$ -regular  $n \times n$  matrix  $A$ . Then  $C$  is equivalent to one of the following 9 matrices.

$$C_1 = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}, C_2 = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}, C_3 = \begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix},$$

$$C_4 = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 \end{bmatrix}, C_5 = \begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{bmatrix}, C_6 = \begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 \end{bmatrix},$$

$C_3^T$ ,  $C_4^T$  and  $C_6^T$ . If  $C = C_4$  then no further columns satisfy equation 4, but then  $\mathbf{c}_2 + \mathbf{c}_3 + \mathbf{c}_4 + \mathbf{c}_5 = \mathbf{j}_5$ , and column  $\mathbf{c}_1$  does not appear, a contradiction. Thus  $C_4$  and  $C_4^T$  can not be submatrices of  $A$ . Similarly,  $C_6$  and  $C_6^T$  can not be submatrices of  $A$ , as the coefficient of  $\mathbf{c}_1$  is negative. Thus every full rank  $5 \times 5$  submatrix has at most three ones in each row/column.

Suppose  $C = C_5$ . Then the only solutions to equation 4 other the columns of  $C_5$  are  $(1, 0, 1, 1, 0)^T$  and  $(0, 0, 0, 1, 0)^T$ . Thus in order to have  $k$  ones in row 5,  $\mathbf{c}_3$  must appear  $k$  times and  $\mathbf{c}_1$  and  $\mathbf{c}_2$  will not appear, a contradiction.

Suppose  $C = C_3$ . The additional solutions to equation 4 are  $\mathbf{c}_6 = \mathbf{c}_1 - \mathbf{c}_4 + \mathbf{c}_5 = (1, 0, 0, 0, 1)^T$ ,  $\mathbf{c}_7 = (0, 0, 1, 0, 0)^T$  and  $\mathbf{c}_8 = (0, 0, 0, 1, 0)^T$ . Since  $[\mathbf{c}_1 \ \mathbf{c}_2 \ \mathbf{c}_3 \ \mathbf{c}_4 \ \mathbf{c}_6]$  is a rank 5 matrix with four ones in row 1,  $\mathbf{c}_6$  does not appear. Then column  $\mathbf{c}_5$  must appear  $k = \frac{n}{4}$  times.

We may now assume that every full rank  $5 \times 5$  submatrix is equivalent to  $C_1$  or  $C_2$ . If  $C = C_2$  then the additional solutions to equation 4 are  $\mathbf{c}_6 = (1, 0, 0, 1, 0)^T$ ,  $\mathbf{c}_7 = (1, 0, 0, 0, 1)^T$  and  $\mathbf{c}_8 = (0, 0, 1, 0, 0)^T$ . Since  $[\mathbf{c}_1 \ \mathbf{c}_2 \ \mathbf{c}_6 \ \mathbf{c}_3 \ \mathbf{c}_5] = C_3$ , column  $\mathbf{c}_6$  does not appear and column  $\mathbf{c}_4$  then appears  $k$  times.

Suppose now that  $C = C_1$ . Additional solutions to equation 4 are  $\mathbf{c}_6 = \mathbf{c}_1 - \mathbf{c}_2 + \mathbf{c}_3 = (0, 1, 1, 0, 0)^T$ ,  $\mathbf{c}_7 = (0, 1, 0, 1, 0)^T$  and  $\mathbf{c}_8 = (0, 1, 0, 0, 1)^T$ . As  $[\mathbf{c}_1 \ \mathbf{c}_6 \ \mathbf{c}_3 \ \mathbf{c}_4 \ \mathbf{c}_5]$  is equivalent to  $C_2$ , we may assume that  $\mathbf{c}_6$  does not appear. Then there must be  $k$  copies of column  $\mathbf{c}_3$ .  $\square$

## 4 Proof of Theorem 1

In this section we will show that there are no directed strongly regular graphs with rank 5,  $t < k$  and with  $\frac{k}{n} \in \{\frac{1}{4}, \frac{2}{7}, \frac{3}{8}, \frac{3}{7}, \frac{4}{9}, \frac{5}{9}, \frac{4}{7}, \frac{5}{8}, \frac{5}{7}, \frac{3}{4}, \frac{4}{5}\}$ . Then the proof of Theorem 1 follows from Theorem 4 and Proposition 3.

Note first that for a dsrg with rank  $r = 5$ , the above mentioned result that  $\frac{k}{n} \leq \frac{2r-3}{2r} = \frac{7}{10}$ , excludes  $\frac{k}{n} = \frac{5}{7}, \frac{3}{4}$ , and  $\frac{4}{5}$ .

**Lemma 11** *If a dsrg with parameters  $(n, k, t, \lambda, \mu)$  has a set  $S$  of vertices, all of which have the same set  $N$  of out-neighbours (or they all have the same set in-neighbours), then*

$$|S| \leq k - \lambda, \quad \text{and} \quad |S| \leq n - 2k + t.$$

**Proof** We have that  $|N| = k$  and  $S \cap N = \emptyset$ . Let  $v \in S$  and  $w \in N$ . Then there are  $\lambda$  paths of length 2 from  $v$  to  $w$ , i.e.,  $w$  has  $\lambda$  in-neighbours in  $N$ . Also every vertex in  $S$  is an in-neighbour of  $w$ . Thus  $|S| + \lambda \leq k$ .

The number of in-neighbours of  $v$  in  $S \cup N$  is  $t$ . Thus  $v$  has  $k - t$  in-neighbours outside  $S \cup N$ . Thus  $|S| + k + (k - t) \leq n$ .  $\square$

**Proposition 12** *A dsrg with rank 5 and  $\frac{k}{n} = \frac{4}{7}$  does not exist.*

**Proof** Suppose that  $(n, k, t, \lambda, \mu)$  are the parameters of a dsrg with rank 5 and with  $\frac{k}{n} = \frac{4}{7}$ . By Proposition 3, we have  $(n, k, t, \lambda, \mu) = (49m, 28m, 20m, 13m, 20m)$ . By Lemma 5, this graph has a set of  $\frac{2}{7} \cdot 49m = 14m$  vertices that either all have the same out-neighbours or all have the same in-neighbours. But this contradicts Lemma 11, as  $n - 2k + t = 13m$ .  $\square$

**Proposition 13** *A dsrg with rank 5 and  $\frac{k}{n} = \frac{1}{4}, \frac{2}{7}, \frac{3}{8}, \frac{4}{9}, \frac{5}{9}$ , or  $\frac{5}{8}$ , does not exist.*

**Proof** In each case we get the parameters from Proposition 3 and apply the first part of Lemma 11.

$\frac{k}{n} = \frac{1}{4}$ : A dsrg with parameters  $(64m, 16m, 5m, m, 5m)$  has  $\frac{1}{4}n = 16m > k - \lambda = 15m$ , a contradiction to Lemma 10.

$\frac{k}{n} = \frac{2}{7}$ : A dsrg with parameters  $(98m, 28m, 10m, 3m, 10m)$  has  $\frac{2}{7}n = 28m > k - \lambda = 25m$ , a contradiction to Lemma 8.

$\frac{k}{n} = \frac{3}{8}$ : A dsrg with parameters  $(256m, 96m, 45m, 21m, 45m)$  has  $\frac{3}{8}n = 96m > k - \lambda = 75m$ , a contradiction to Lemma 7.

$\frac{k}{n} = \frac{4}{9}$ : A dsrg with parameters  $(81m, 36m, 20m, 19m, 20m)$  has  $\frac{4}{9}n = 36m > k - \lambda = 17m$ , a contradiction to Lemma 9.

$\frac{k}{n} = \frac{5}{9}$ : A dsrg with parameters  $(324m, 180m, 125m, 90m, 125m)$  has  $\frac{5}{9}n = 180m > k - \lambda = 90m$ , a contradiction to Lemma 9.

$\frac{k}{n} = \frac{5}{8}$ : A dsrg with parameters  $(256m, 160m, 125m, 85m, 125m)$  has  $\frac{5}{8}n = 160m > k - \lambda = 75m$ , a contradiction to Lemma 7.  $\square$

**Proposition 14** *A dsrg with rank 5 and  $\frac{k}{n} = \frac{3}{7}$  does not exist.*

**Proof** Suppose that  $(n, k, t, \lambda, \mu)$  are the parameters of a dsrg with rank 5 and with  $\frac{k}{n} = \frac{3}{7}$ . By Proposition 3, we have  $(n, k, t, \lambda, \mu) = (196m, 84m, 45m, 24m, 45m)$ . Let  $A$  be the adjacency matrix of this dsrg. Then  $A^T$  is the adjacency matrix of a dsrg with the same parameters. By Lemma 11, a set  $S$  of vertices with identical out- (or in-) neighbour sets has  $|S| \leq k - \lambda = 60m < \frac{3}{7}n$ . Therefore by Corollary 6, using  $A^T$  if necessary, we have two disjoint sets  $S_1$  and  $S_2$  of vertices with  $|S_1| = |S_2| = \frac{2}{7}n = 56m$  and sets  $N_1$  and  $N_2$  with  $|N_1| = |N_2| = 84m$  so that every vertex in  $S_i$  has out-neighbour set  $N_i$ . Then  $N_1 \cap N_2 = \emptyset$ , as a vertex in  $N_1 \cap N_2$  would have in-degree at least  $|S_1 \cup S_2| = 112m$ . We also have  $S_i \cap N_i = \emptyset$ , but since there are only  $28m$  vertices outside  $N_1 \cup N_2$ , the sets  $S_1 \cap N_2$  and  $S_2 \cap N_1$  are non-empty. Let  $v \in S_1 \cap N_2$ . A subset of  $N_1$  of exactly  $t = 45m$  vertices are in-neighbours of  $v$ .  $S_2 \cap N_1$  is contained in this set. Thus  $|S_2 \cap N_1| \leq 45m$ , and so there exists a vertex  $x \in S_2 \setminus N_1$ . Similarly there exists  $y \in S_1 \setminus N_2$ . Since  $x$  and  $y$  are non-adjacent there are  $45m$  paths from  $x$  to  $y$  of length 2. The internal vertex of such a path belongs to  $N_2 \setminus S_1$ , as  $S_1$  is independent. Thus  $|N_2 \setminus S_1| \geq 45m$  and so  $|S_1 \setminus N_2| \geq 17m$ . Similarly,  $|S_2 \setminus N_1| \geq 17m$ . But these sets are disjoint subsets of a set of  $28m$  vertices, a contradiction.  $\square$

## References

- [1] A. Araluze, I. Kovács, K. Kutnar, L. Martínez, D. Marušič, Partial sum quadruples and bi-Abelian digraphs, *Journal of Combinatorial Theory (A)* **119** (2012) 1811–1831.
- [2] A.E. Brouwer, S.A. Hobart, Parameters of directed strongly regular graphs, <http://homepages.cwi.nl/~aeb/math/dsrg/dsrg.html>.
- [3] A. E. Brouwer, O. Olmez and S. Y. Song, Directed strongly regular graphs from  $1\frac{1}{2}$ -designs, *European Journal of Combinatorics* **33** (2012) 1174–1177.
- [4] A. M. Duval, A Directed Version of Strongly Regular Graphs, *Journal of Combinatorial Theory (A)* **47** (1988) 71–100.
- [5] F. Fiedler, M. H. Klin, M. Muzychuk, Small vertex-transitive directed strongly regular graphs, *Discrete Mathematics* **255** (2002) 87–115.

- [6] C. D. Godsil, S. A. Hobart and W. J. Martin, Representations of directed strongly regular graphs, *European Journal of Combinatorics* **28** (2007) 1980–1993.
- [7] L. K. Jørgensen, Directed strongly regular graphs with  $\mu = \lambda$ , *Discrete Mathematics* **231** (2001) 289–293.
- [8] L. K. Jørgensen, Non-existence of directed strongly regular graphs, *Discrete Mathematics* **264** (2003) 111–126.
- [9] L. K. Jørgensen, Rank of adjacency matrices of directed (strongly) regular graphs, *Linear Algebra and its Applications* **407** (2005) 233–241.
- [10] M. Klin, A. Munemasa, M. Muzychuk and P.-H. Zieschang, Directed strongly regular graphs obtained from coherent algebras, *Linear Algebra and its Applications* **377** (2004) 83–109.